

CONF-850132--1

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

LA-UR--84-3447

DEG5 002434

**TITLE** The Structure and Forcing of Teleconnection Patterns  
Occurring in General Circulation Model Simulations

**AUTHOR(S)** Robert C. Malone, Los Alamos National Laboratory  
Eric J. Pitcher, University of Miami  
Maurice L. Blackmon, National Center for Atmospheric Research  
John E. Geisler, University of Utah

**SUBMITTED TO** Third Conference on Climate Variations  
Los Angeles, CA  
January 8-11, 1985

#### DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

**Los Alamos** Los Alamos National Laboratory  
Los Alamos, New Mexico 87545

FORM NO 836 R4  
ST NO 8070 5/81

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

*JSW*

THE STRUCTURE AND FORCING OF TELECONNECTION PATTERNS  
OCCURRING IN GENERAL CIRCULATION MODEL SIMULATIONS

Robert C. Malone  
Los Alamos National Laboratory, Los Alamos, New Mexico

Eric J. Pitcher  
University of Miami, Miami, Florida

Maurice L. Blackmon  
National Center for Atmospheric Research, Boulder, Colorado

John E. Geisler  
University of Utah, Salt Lake City, Utah

Several physical mechanisms have been proposed to explain the occurrence of midlatitude atmospheric teleconnection patterns. We consider here two such mechanisms, Rossby wave trains forced by tropical heating (Hoskins and Karoly, 1981; Simmons, 1982) and the growth of barotropically unstable modes in midlatitudes (Simmons *et al.*, 1983). Because these mechanisms have been discussed in the literature in essentially steady-state form, and because both mechanisms appear capable of explaining the steady-state aspects of observed teleconnection patterns, we consider instead features associated with presumed time-dependent characteristics of these two mechanisms. The two mechanisms appear to have quite different time-dependent characteristics.

We examine several characteristics of teleconnection-pattern fluctuations occurring in GCM simulations in an attempt to discriminate between these two proposed mechanisms. The model used is the NCAR Community Climate Model, whose simulation characteristics are discussed in Pitcher *et al.* (1983), Ramanathan *et al.* (1983) and Malone *et al.* (1984). Five model simulations are considered: a perpetual-January control case with climatological-mean sea-surface temperatures (SST) and four cases in which SST anomalies of various intensities have been introduced in the tropical Pacific. The SST anomaly pattern is that used by Blackmon *et al.* (1983) in their study of changes in the time-mean state of the atmosphere associated with equatorial Pacific SST anomalies. For brevity, we restrict our attention to the model's Pacific-North American (PNA) teleconnection pattern.

Our findings may be summarized as follows. (1) We find that teleconnection fluctuations with the PNA pattern appear in all five simulations and that the amplitude of the PNA-pattern fluctuations (computed as fluctuations relative to the time-mean of each case individually) is relatively insensitive to changes in the mean tropical SST forcing. (2) Although each SST distribution is fixed in time, the heating rate due to convective processes in the tropical atmosphere can fluctuate. We have tested the hypothesis that tropical heating fluctuations directly force midlatitude geopotential-height fluctuations by

performing a cross-spectral analysis of the fluctuations in these two fields, and find no evidence for direct forcing. This is not to be confused with tropical forcing of anomalies in the time-averaged extratropical height field, which Blackmon *et al.* (1983) have shown does occur in these same simulations. (3) In this same analysis, we did find a highly significant correlation (with a 30-day period) between height fluctuations near 47°N, 165°W and subtropical convective heating to the southeast near 20°N, 135°W; the height fluctuations are in phase with or slightly lead the heating fluctuations. (4) By means of a cross-spectral analysis of height fluctuations at the centers of action (antinodes) of the PNA pattern and at the "nodal" points in between, we conclude that the oscillations linking the Tropical Pacific (TP) and North Pacific (NP) points constitute pure standing oscillations having very high coherence for all periods of 20 days or longer (Figs. 1 and 2). There is no evidence for propagation between these two points. The oscillations connecting the NP and North American (NA) points are of mixed character, appearing as traveling waves at some frequencies and not at others (Fig. 3).

The apparent dependence on geographical position of the character of the oscillation (standing, traveling or indeterminate) is quite puzzling. However similar features can be seen in the complex phase structure of the unstable barotropic normal mode displayed in Fig. 12 of Simmons *et al.* (1983). In particular, the geopotential height in the tropical central Pacific oscillates out of phase with that in the north Pacific and a distinct node lies between these regions. This mode also has a period of about 30 days, which suggests a possible explanation for the appearance of this period in the simulation analyses.

While the evidence is certainly not conclusive, it suggests that the short term (one to several months) fluctuations in the PNA pattern are associated with the unstable barotropic normal mode. Heating by tropical SST anomalies appears most influential when averages are taken over a season or longer.




Fig. 1. Coherence and phase from cross-spectral analyses of the 500-mb height fluctuations at TP and NP, at NP and NA, and also at TP with the 500-mb convective heating rate fluctuations in an area directly south of the TP point. Significance levels are indicated on the right side.




Fig. 2. As in Fig. 1, but for the fluctuations at both TP and NP with those at the intermediate nodal point. The (TP, NP) curve is repeated from Fig. 1.




Fig. 3. As in Fig. 1, but for the fluctuations at both NP and NA with those at the intermediate nodal point. Twice as many lags were used in this analysis as were used in Figs. 1 and 2.

#### References

- Blackmon, M. L., J. E. Geisler, and E. J. Pitcher, 1983: A general circulation model study of January climate anomaly patterns associated with interannual variation of equatorial Pacific sea surface temperatures. *J. Atmos. Sci.*, **40**, 1410-1425.
- Brankin, E. J. and D. J. Karoly, 1981: The steady linear response of a spherical atmosphere to thermal and orographic forcing. *J. Atmos. Sci.*, **38**, 1179-1196.
- Malone, R. C., E. J. Pitcher, M. L. Blackmon, K. Puri and W. Bourke, 1984: The simulation of stationary and transient geopotential-height eddies in January and July with a spectral general circulation model. *J. Atmos. Sci.*, **41**, 1394-1419.
- Pitcher, E. J., R. C. Malone, V. Ramanathan, M. L. Blackmon, K. Puri and W. Bourke, 1983: January and July simulations with a spectral general circulation model. *J. Atmos. Sci.*, **40**, 580-604.
- Ramanathan, V., E. J. Pitcher, R. C. Malone and M. L. Blackmon, 1983: The response of a spectral general circulation model to refinements in radiative processes. *J. Atmos. Sci.*, **40**, 605-630.
- Simmons, A. J., 1982: The forcing of stationary wave motion by tropical diabatic heating. *Quart. J. Roy. Meteor. Soc.*, **108**, 503-534.
- Simmons, A. J., J. M. Wallace, and G. W. Branstator, 1983: Barotropic Wave Propagation and Instability, and Atmospheric Teleconnection Patterns. *J. Atmos. Sci.*, **40**, 1363-1392.